

# 注射對比劑同意書

## Radiology Department of Taichung Armed Forces General Hospital

### Contrast Agent Injection Consent Form

Examinations involving the intravenous injection of a contrast agent utilize the contrast's effect on the area to be examined to enhance and improve the visibility of the organs or lesions under examination to reach a diagnosis. Conventional ionic iodine-based contrast agents may cause allergic reactions in a small number of patients (the incidence of allergic reactions is 10%). Symptoms include nausea, vomiting, skin rashes, itching, and in a few severe cases, shock or even sudden death (the reported mortality rate is approximately 1 in 100,000, according to international statistics). However, if you have any of the following conditions, you have a significantly increased risk of allergy.

- ☐ 1. Heart, lung, and vascular diseases (impaired cardiac function, hypertension, arteriosclerosis and emphysema).
- ☐ 2. Cirrhosis, liver dysfunction.
- ☐ 3. Kidney dysfunction, uremia.
- ☐ 4. Asthma.
- ☐ 5. Diabetes mellitus for five years or longer.
- ☐ 6. History of allergies (including reactions caused by the injection of iodine-based contrast agents, and allergies to food, other medications, pollen, or other allergens).
- ☐ 7. A senior age with poor physical condition (72 years or above).
- ☐ 8. A young age (infants or small children of 2 years or younger).
- ☐ 9. Critical illness.

If you have any of the above conditions, please indicate a ~ in the corresponding ☐

Signature of patient or family: \_\_\_\_\_

If the patient is a minor, a legal representative should sign the consent form.

※Please read these instructions before deciding to sign the examination consent form※

Date:                      YYYY                      MM                      DD                      Time:                      hh                      mm